arXiv:1408.6727v1 [math.PR] 28 Aug 2014

EXACT DISTRIBUTION OF VERHULST PROCESS

JACEK JAKUBOWSKI AND MACIEJ WIŚNIEWOLSKI

Institute of Mathematics, University of Warsaw
Banacha 2, 02-097 Warszawa, Poland
e-mail: jakub@mimuw.edu.pl
and
wisniewolski@mimuw.edu.pl

Abstract. We investigate a Verhulst process, which is the special functional of geometric Brownian motion and has many applications, among others in biology and in stochastic volatility models. We present an exact form of density of a one dimensional distribution of Verhulst process. Simple formula for the density of Verhulst process is obtained in the special case, when the drift of geometric Brownian motion is equal to $-\frac{1}{2}$. Some special properties of this process are discussed, e.g. it turns out that under Girsanov's change of measure a Verhulst process still remains a Verhulst process but with different parameters.

Key words: geometric Brownian motion, Verhulst process, Girsanov's change of measure, Laplace transform, exponential functional of Brownian motion

2010 AMS Subject Classification: 60J65, 60J70.

1. Introduction

The paper is devoted to study the distribution of Verhulst process. A Verhulst process is the special functional of geometric Brownian motion with drift and its integral. The process had been studied among others by Mackevičius [7] and Lungu and Øksendal [6], where it is called a process of population growth in stochastic crowded environment. The results in both papers were obtained by a some kind of approximations. In Øksendal [14] there were indicated some applications of a Verhulst process in finance and biology. Mackevičius [8] during his conference presentation on Vilnius Conference 2014 indicates on the need to find the closed-form expressions for Laplace transform and exact distribution of Verhulst process. In this paper we provide the formulas for one dimensional distribution of the process. Results obtained by Yor

and collaborators in several papers and monographes (see, e.g. [2], [3], [4], [9], [11], [12], [13]) on the distribution of $(B_t^{(\mu)}, \int_0^t e^{B_u^{(\mu)}} du)$, where $B_t^{(\mu)} = B_t + \mu t$ is a Brownian motion with drift, give us the background in providing closed formulas for the density of Verhulst process, which in case of drift $\mu = -\frac{1}{2}$ becomes especially simple. We present also some interesting and important properties of the Verhulst process among them the fact that a Verhulst process remains, under Girsanov's change of measure, a Verhulst proces but with different parameters. The ideas presented below are original and are not easy consequences of the previous results.

2. Distribution and properties of Verhulst process

We work on a complete probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with filtration $(\mathcal{F}_t)_{t\in[0,\infty)}$ and Brownian motion B defined on it. We define a Verhulst process $\theta^{(\mu,\beta)}$ starting from 1 as the functional of a Brownian motion with drift

$$\theta_t^{(\mu,\beta)} = \frac{e^{B_t+\mu t}}{1+\beta\int_0^t e^{B_s+\mu s}ds}. \tag{1}$$

It is easy to see that this functional is the unique strong solution of SDE

$$d\theta_t^{(\mu,\beta)} = \theta_t^{(\mu,\beta)}dB_t + \Big((\mu+1/2)\theta_t^{(\mu,\beta)} - \beta(\theta_t^{(\mu,\beta)})^2\Big)dt,\ \theta_0^{(\mu,\beta)} = 1, \tag{2}$$

for $\mu \in \mathbb{R}$, $\beta \geq 0$ and $t \geq 0$. It is worth to note that taking $\theta_t^{(\mu,\beta)}$ starting from any $x > 0$ does not change any of its probabilistic properties (see also Section 3). As it was mentioned, the process $\theta^{(\mu,\beta)}$ is also known in literature as a process of population growth in crowded stochastic environment process (see [14]). The approximation of the process was presented in [7]. The properties of Verhulst process were also studied by Jakubowski and Wiśniewolski [5]. In particular it was shown there (see Theorem 2.19) that for $\lambda > 0, a \in \mathbb{R}$, we have

$$\mathbb{E}\Big[\exp\Big(-\frac{\lambda e^{aB_t^{(\mu)}}}{1+\beta A_t^{(\mu)}}\Big)\Big] = \mathbb{E}\Big[F_{B_t^{(\mu)}}\Big(\beta^{-1}R^{(\lambda e^{aB_t^{(\mu)}})}(1/2)\Big)\Big], \tag{3}$$

where $B_t^{(\mu)} = B_t + \mu t$, $A_t^{(\mu)} = \int_0^t e^{2B_s^{(\mu)}}ds$, and F_x is given by

$$F_x(z) = \exp\Big(-\frac{\varphi_z(x)^2 - x^2}{2t}\Big) \tag{4}$$

with

$$\varphi_x(y) = \operatorname{arcosh}(xe^{-y} + \cosh(y)) \tag{5}$$
$$= \ln\Big(xe^{-y} + \cosh(y) + \sqrt{x^2e^{-2y} + \sinh^2(y) + 2xe^{-y}\cosh(y)}\Big).$$

Moreover, R^x is a squared Bessel process with the index -1 starting at x and independent of $(B_t^{(\mu)}, t \geq 0)$.

Observe that from (3) we easily obtain the Laplace transform of $\theta_t^{(\mu,\beta)}$.

Proposition 2.1. *For $\lambda > 0$ we have*

$$\mathbb{E}e^{-\lambda\theta_t^{(\mu,\beta)}} = \mathbb{E}\Big[F_{B_t^{(2\mu)}}\Big((4\beta)^{-1}R^{(\lambda e^{2B_t^{(2\mu)}})}(1/2)\Big)\Big]. \tag{6}$$

Proof. Exactly in the same manner as in the proof of Theorem 2.20 in [5], we have

$$\theta_{4t}^{(\mu,\beta)} = \frac{e^{2(B_{4t}/2+2t\mu)}}{1 + 4\beta\int_0^t e^{2(B_{4t}/2+2t\mu)}du}.$$

Since $B_{4t}/2$ is a Brownian motion, the statement follows from (3) with $a = 2$, μ replaced by 2μ and β replaced by 4β. □

The following lemma will be used further.

Lemma 2.2. *Fix $\gamma > 0$, $\mu \in \mathbb{R}$, $\beta \geq 0$. Then*

$$M_t^{(\gamma,\mu,\beta)} = e^{-\gamma\int_0^t \theta_s^{(\mu,\beta)}dB_s - \frac{\gamma^2}{2}\int_0^t(\theta_s^{(\mu,\beta)})^2ds}, \qquad t \in [0,\infty) \tag{7}$$

is a martingale.

Proof. It is enough to prove that for fixed $T > 0$ the local martingale $M_t^{(\gamma,\mu,\beta)}, t \in [0,T]$ is bounded. It is so, since SDE (2) implies that

$$\exp\Big(-\gamma\int_0^t \theta_s^{(\mu,\beta)}dB_s - \frac{\gamma^2}{2}\int_0^t(\theta_s^{(\mu,\beta)})^2ds\Big)$$
$$= e^{-\gamma\left(\theta_t^{(\mu,\beta)} - 1 - (\mu+1/2)\int_0^t\theta_s^{(\mu,\beta)}ds + \beta\int_0^t(\theta_s^{(\mu,\beta)})^2ds\right) - \frac{\gamma^2}{2}\int_0^t(\theta_s^{(\mu,\beta)})^2ds}$$
$$= e^{-\gamma(\theta_t^{(\mu,\beta)}-1) - (\gamma\beta+\frac{\gamma^2}{2})\int_0^t\left((\theta_s^{(\mu,\beta)})^2 - 2\theta_s^{(\mu,\beta)}\frac{\gamma(\mu+1/2)}{\gamma^2+2\gamma\beta} + \left(\frac{\gamma(\mu+1/2)}{\gamma^2+2\gamma\beta}\right)^2\right)ds} \times$$
$$\times\, e^{\frac{(\gamma(\mu+1/2))^2}{4(\gamma\beta+\gamma^2/2)}t} < \infty.$$

□

Remark 2.3. One can wonder if $\overline{M}_t^{(\gamma,\mu,\beta)} = e^{\gamma\int_0^t\theta_s^{(\mu,\beta)}dB_s - \frac{\gamma^2}{2}\int_0^t(\theta_s^{(\mu,\beta)})^2ds}$, for a fixed $\gamma > 0$, could be a martingale as well. In Remark 1.1 [4] it was noticed that $\overline{M}^{(\gamma,\mu,\beta)}$ can not be a martingale.

Lemma 2.2 enables us to define a new probability measure

$$\frac{d\mathbb{Q}^{(\gamma,\mu,\beta,T)}}{d\mathbb{P}}\Big|_{\mathcal{F}_T} = M_T^{(\gamma,\mu,\beta)}. \tag{8}$$

From Girsanov's theorem $V_s = B_s + \gamma \int_0^s \theta_u^{(\mu,\beta)} du$, $s \le T$, is a Brownian motion under $\mathbb{Q}^{(\gamma,\mu,\beta)}$. This leads us to the following

Theorem 2.4. *Let $\theta_t^{(\mu,\beta)}$ be defined by (1) and $\mathbb{Q}^{(\gamma,\mu,\beta,T)}$ by (8). Then*

$$\theta_t^{(\mu,\beta)} = \frac{e^{V_t+\mu t}}{1+(\beta+\gamma)\int_0^t e^{V_u+\mu u}du}, \qquad t \le T, \tag{9}$$

where V is a standard Brownian motion under $\mathbb{Q}^{(\gamma,\mu,\beta,T)}$.

Proof. From Ito's lemma and (2) follows that

$$\ln \theta_t^{(\mu,\beta)} = B_t + \int_0^t (\mu - \beta\theta_s^{(\mu,\beta)})ds. \tag{10}$$

Taking $V_t = B_t + \gamma \int_0^t \theta_u^{(\mu,\beta)} du$, a Brownian motion under $\mathbb{Q}^{(\gamma,\mu,\beta,T)}$, we obtain from the last equality that

$$V_t + \mu t = \ln \theta_t^{(\mu,\beta)} + (\beta+\gamma)\int_0^t \theta_u^{(\mu,\beta)} du. \tag{11}$$

Thus direct computation yields

$$\begin{aligned}\int_0^t e^{V_u+\mu u}du &= \int_0^t \theta_u^{(\mu,\beta)} e^{(\beta+\gamma)\int_0^u \theta_s^{(\mu,\beta)}ds}du \\ &= \frac{1}{\gamma+\beta}\Big(e^{(\beta+\gamma)\int_0^t \theta_u^{(\mu,\beta)}du} - 1\Big).\end{aligned}$$

From the last equality and (11) we finally have

$$\theta_t^{(\mu,\beta)} = \frac{e^{V_t+\mu t}}{1+(\beta+\gamma)\int_0^t e^{V_u+\mu u}du}.$$

□

This theorem justifies that a Verhulst process $\theta^{(\mu,\beta)}$ on $(\Omega,\mathcal{F},\mathbb{P})$ remains a Verhulst process under $\mathbb{Q}^{(\gamma,\mu,\beta,T)}$, though with different parameters.

Proposition 2.5. *If $(\theta_t^{(\mu,\beta)})$, $t \le T$, is a Verhulst process with parameters (μ,β) on $(\Omega,\mathcal{F},\mathbb{P})$, then $(\theta_t^{(\mu,\beta)})$ is on $(\Omega,\mathcal{F},\mathbb{Q}^{(\gamma,\mu,\beta,T)})$ a Verhulst process with parameters $(\mu,\gamma+\beta)$.*

We are ready to derive the formula for the density of one dimensional distribution of Verhulst process, i.e. the density of $\theta_t^{(\mu,\beta)}$ for a fixed $t > 0$. Let us introduce the following notation

$$\mathbb{P}(\theta_t^{(\mu,\beta)} \in dx) = g_t(\beta, x)dx.$$

Observe that for $\beta = 0$, $g_t(0, x)$ is the density of geometric Brownian motion $e^{B_t+\mu t}$. Proposition 2.5 implies that

$$g_t(\beta + \gamma, x) = g_t(\beta, x)\mathbb{E}(M_t^{(\gamma,\mu,\beta)}|\theta_t^{(\mu,\beta)} = x).$$

In particular, for $\beta = 0$ we obtain

$$g_t(\gamma, x) = g_t(0, x)\mathbb{E}(M_t^{(\gamma,\mu,0)}|e^{B_t+\mu t} = x). \tag{12}$$

Theorem 8.1 in Matsumoto - Yor [12] states that:
for any $t > 0, \lambda > 0, v > 0, x \in \mathbb{R}$

$$\begin{aligned}\psi_t^{(\mu)}(v, x)&\mathbb{E}\Big(e^{-\frac{\lambda^2}{2}\int_0^t e^{2B_u+2\mu u}du}\Big|\int_0^t e^{B_u+\mu u}du = v, B_t + \mu t = x\Big) \\ &= e^{\mu x-\mu^2 t/2}\frac{\lambda}{4\sinh(\lambda v/2)}e^{-\lambda(1+e^x)\coth(\lambda v/2)}\Theta(\phi(v, x, \lambda), t/4),\end{aligned} \tag{13}$$

where

$$\begin{aligned}\psi_t^{(\mu)}(v, x) &= \frac{1}{16}e^{\mu x-\mu^2 t/2}\frac{1}{v}e^{-\frac{2(1+e^x)}{v}}\Theta(4e^{x/2}/v, t/4), \\ \Theta(r, t) &= \frac{r}{\sqrt{2\pi^3 t}}e^{\frac{\pi^2}{2t}}\int_0^\infty e^{-\frac{z^2}{2t}}e^{-r\cos(z)}\sinh(z)\sin(\pi z/t)dz, \\ \phi(v, x, \lambda) &= \frac{2\lambda e^{x/2}}{\sinh(\lambda v/2)}.\end{aligned}$$

Using this result we can write the formula for density of Verhulst process.

Theorem 2.6. *The density of Verhulst process is given by*

$$g_t(\gamma, x) = g_t(0, x)e^{-\gamma(x-1)}\mathbb{E}H_t(a_t^{(\mu)}, x),$$

where $a_t^{(\mu)} = \int_0^t e^{B_u+\mu u}du$, *and*

$$H_t(y, x) = e^{\gamma(\mu+1/2)y}\mathbb{E}\Big(e^{-\frac{\gamma^2}{2}\int_0^t e^{2B_u+2\mu u}du}|a_t^{(\mu)} = y, B_t + \mu t = \ln x\Big).$$

Proof. We have from (7) and (2)

$$\begin{aligned}\mathbb{E}(M_t^{(\gamma,\mu,0)}|e^{B_t+\mu t}=x) &= e^{-\gamma(x-1)}\mathbb{E}\Big(e^{\gamma(\mu+1/2)a_t^{(\mu)}-\frac{\gamma^2}{2}\int_0^t e^{2B_u+2\mu u}du}|e^{B_t+\mu t}=x\Big)\\ &= e^{-\gamma(x-1)}\int_0^\infty H_t(y,x)\mathbb{P}(a_t^{(\mu)}\in dy)\\ &= e^{-\gamma(x-1)}\mathbb{E}H_t(a_t^{(\mu)},x).\end{aligned}$$

Thus, the result follows from (12). □

Remark 2.7. Taking together Theorem 2.6 and (13) enable us to write down the close formula for density of Verhulst process. The density of $a_t^{(\mu)}$ can be obtained from the formula

$$\mathbb{P}(a_t^{(\mu)}\in dv, B_t+\mu t\in dx)=\psi_t^{(\mu)}(v,x)dvdx$$

(see [12, Theorem 4.1]). Another formula for density of $a_t^{(\mu)}$ can be also found on page 612 in [1], formula (1.8.4).

Proposition 2.8. *For a Verhulst process $\theta_t^{(\mu,\beta)}$, a Brownian motion B, $\beta>0$ and $\mu\neq -1/2$ we have*

$$\begin{aligned}\theta_t^{(\mu,\beta)}e^{\beta\int_0^t\theta_s^{(\mu,\beta)}ds} &= e^{B_t+\mu t},\\ \mathbb{E}e^{\beta\int_0^t\theta_s^{(\mu,\beta)}ds} &= 1+\frac{\beta}{\mu+1/2}\Big(e^{(\mu+1/2)t}-1\Big).\end{aligned}$$

Proof. The first equality follows immediately from (10). For the second observe that

$$e^{\beta\int_0^t\theta_s^{(\mu,\beta)}ds}=(1+\beta\int_0^t e^{B_u+\mu u}du), \tag{14}$$

so

$$e^{(\mu+1/2)t}=\mathbb{E}e^{B_t+\mu t}=\mathbb{E}\theta_t^{(\mu,\beta)}e^{\beta\int_0^t\theta_s^{(\mu,\beta)}ds}=\frac{1}{\beta}\frac{\partial}{\partial t}\mathbb{E}e^{\beta\int_0^t\theta_s^{(\mu,\beta)}ds}.$$

Thus

$$\mathbb{E}e^{\beta\int_0^t\theta_s^{(\mu,\beta)}ds}=1+\frac{\beta}{\mu+1/2}\Big(e^{(\mu+1/2)t}-1\Big).$$

□

As an example of applications we present a solution to problem of finding a special representation of Brownian motion with drift. For another distributional equations of this kind see e.g. Section 13 in [10].

Example 2.9. Let B be a Brownian motion under $\mathbb{P}$, $\mu \in \mathbb{R}$ and $\alpha \in (0,1)$. Our aim is to find a measure $\mathbb{Q}$, a Brownian motion V under $\mathbb{Q}$ and a random variable G such that distribution of G under $\mathbb{P}$ and under $\mathbb{Q}$ belongs to the same class, and for fixed $t > 0$

$$B_t + \mu t = \alpha(V_t + \mu t) + (1-\alpha)G. \tag{15}$$

Fix T, $T > t$, and $\gamma > 0$. To find a representation (15) we take $\mathbb{Q} = \mathbb{Q}^{(\gamma,\mu,\beta,T)}$ given by (8) with $\beta = \frac{\gamma\alpha}{1-\alpha}$. Then $V_t = B_t + \gamma \int_0^t \theta_s^{(\mu,\beta)} ds$ is a Brownian motion under $\mathbb{Q}$. By (10) and (14) we have

$$e^{V_t+\mu t} = \theta_t^{(\mu,\beta)} \Big(1 + \beta \int_0^t e^{B_s+\mu s} ds\Big)^{\frac{\beta+\gamma}{\beta}}.$$

From the definition of $\theta_t^{(\mu,\beta)}$ and the last equality we obtain

$$e^{B_t+\mu t} = e^{\frac{\beta}{\beta+\gamma}(V_t+\mu t)} (\theta_t^{(\mu,\beta)})^{\frac{\gamma}{\beta+\gamma}}. \tag{16}$$

Since $\alpha = \frac{\beta}{\beta+\gamma}$, by taking ln of both sides of (16) we have

$$B_t + \mu t = \alpha(V_t + \mu t) + (1-\alpha) \ln \theta_t^{(\mu,\beta)},$$

which is (15) with $G = \ln \theta_t^{(\mu,\beta)}$. Observe that $\ln \theta_t^{(\mu,\beta)}$ under both $\mathbb{P}$ and $\mathbb{Q}$ belongs to the same family of ln of Verhulst process. This finishes the proof.

Now, we present another formula for Laplace transform of Verhulst process.

Proposition 2.10. *Let $\theta_t^{(\mu,\beta)}$ be a Verhulst process with parameters (μ,β). Then for $\lambda \geq 0$*

$$\mathbb{E}e^{-\lambda\theta_t^{(\mu,\beta)}} = e^{\beta} \mathbb{E} e^{-(\beta+\lambda)e^{B_t+\mu t} + \beta(\mu+1/2)\int_0^t e^{B_s+\mu s} ds - \frac{\beta^2}{2}\int_0^t e^{2B_s+2\mu s} ds}, \tag{17}$$

where B is a standard Brownian motion under $\mathbb{P}$.

Proof. From Proposition 2.5 we know that a geometric Brownian motion $e^{B_t+\mu t}$ under $\mathbb{P}$ becomes a Verhulst process $\tilde{\theta}_t^{(\mu,\beta)}$ with parameters (μ,β) under $\mathbb{Q} = \mathbb{Q}^{(\beta,\mu,0,t)}$ given by (8). Thus

$$\mathbb{E}e^{-\lambda\theta_t^{(\mu,\beta)}} = \mathbb{E}_{\mathbb{Q}} e^{-\lambda\tilde{\theta}_t^{(\mu,\beta)}} = \mathbb{E} e^{-\lambda e^{B_t+\mu t}} M_t^{(\beta,\mu,0)},$$

where $M^{(\beta,\mu,0)}$ is defined by (7). The assertion follows from (2) and some simple algebra. □

3. Exponential random time and drift $\mu = -\frac{1}{2}$

In this section we will consider a Verhulst process θ starting from $x > 0$ with $\mu = -\frac{1}{2}$ and $\beta = x$. Thus, θ is of the special form

$$\theta_t = \frac{xe^{B_t - \frac{t}{2}}}{1 + x\int_0^t e^{B_u - \frac{u}{2}} du}, \tag{18}$$

where B is a Brownian motion. Let T_λ be an exponential random variable with parameter $\lambda > 0$, independent of B. We have

Lemma 3.1. *Let $v = \sqrt{2\lambda + 1/4}$. The density of θ_{T_λ} is given on $(0, \infty)$ by*

$$\mathbb{P}(\theta_{T_\lambda} \in dz) = 2\lambda e^{x-z}\sqrt{x/z^3} F_v(x, z)dz, \tag{19}$$

where

$$F_v(x, y) = I_v(x \vee y)K_v(x \wedge y)$$

is a product of two modified Bessel functions.

Proof. For $r \geq 0$ from Proposition 2.10 (where we put $\mu = -1/2, \lambda = xr, \gamma = x$) we obtain

$$\mathbb{E}e^{-r\theta_t} = e^x \mathbb{E}e^{-x(r+1)e^{B_t - t/2} - \frac{x^2}{2}\int_0^t e^{2B_u - u} du}.$$

Thus, using [12, Theorem 4.11] describing the joint density of the vector $(e^{B_{T_\lambda} - T_\lambda/2}, \int_0^{T_\lambda} e^{2B_u - u} du)$, we have

$$\begin{aligned} \mathbb{E}e^{-r\theta_{T_\lambda}} &= e^x \mathbb{E}e^{-x(r+1)e^{B_{T_\lambda} - T_\lambda/2} - \frac{x^2}{2}\int_0^{T_\lambda} e^{2B_u - u} du} \\ &= e^x \int_0^\infty \int_0^\infty e^{-x(r+1)y - \frac{x^2}{2}w} \frac{\lambda}{y^{v+5/2}} p^{(v)}(w, 1, y)dydw, \end{aligned} \tag{20}$$

where $p^{(v)}$ is the transition probability density of the Bessel process with index v. Again from [12] (see Remark 2.1) we have for $\alpha > 0$

$$\int_0^\infty e^{-\alpha t} p^{(v)}(t, x, y)dt = 2y\Big(\frac{y}{x}\Big)^v F_v(\sqrt{2\alpha}x, \sqrt{2\alpha}y).$$

Thus by (20), Fubbini's theorem and the last identity we have

$$\begin{aligned} & e^x \int_0^\infty \int_0^\infty e^{-x(r+1)y - \frac{x^2}{2}w} \frac{\lambda}{y^{v+5/2}} p^{(v)}(w, 1, y)dydw \\ &= \int_0^\infty e^{x - x(r+1)y} \frac{\lambda}{y^{v+5/2}} 2y^{v+1} F_v(x, xy)dy \\ &= \int_0^\infty e^{x - rw - w} \frac{2\lambda}{w^{3/2}} \sqrt{x} F_v(x, w)dw \end{aligned}$$

and the assertion follows from the standard Laplace transform argument. □

Now we are ready to derive the exact formula for the density of θ_t.

Theorem 3.2. *Fix $t > 0$. Then on $(0, \infty)$*

$$\mathbb{P}(\theta_t \in dw) = e^{-\frac{t}{8}+x-w}\sqrt{\frac{x}{w^3}}\Big(\int_0^\infty \frac{1}{z}e^{-\frac{z}{2}-\frac{x^2+w^2}{2z}}\Theta(xw/z,t)dz\Big)dw, \tag{21}$$

where

$$\Theta(r,t) = \frac{r}{\sqrt{2\pi^3 t}}e^{\frac{\pi^2}{2t}}\int_0^\infty e^{-\frac{z^2}{2t}}e^{-r\cos(z)}\sinh(z)\sin(\pi z/t)dz. \tag{22}$$

Proof. We have on $(0, \infty)$

$$\mathbb{P}(\theta_{T_\lambda} \in dw) = \lambda\int_0^\infty e^{-\lambda t}\mathbb{P}(\theta_t \in dw)dt. \tag{23}$$

where T_λ is an exponential random variable with parameter λ independent of the process θ. On the other side, from Lemma 3.1 for $\lambda > 0$ and $v = \sqrt{2\lambda + 1/4}$, we have on $(0, \infty)$

$$\begin{aligned}
\mathbb{P}(\theta_{T_\lambda} \in dw) &= 2\lambda e^{x-w}\sqrt{\frac{x}{w^3}}F_v(x,w)dw \\
&= 2\lambda e^{x-w}\sqrt{\frac{x}{w^3}}I_v(x\vee w)K_v(x\wedge w)dw \\
&= 2\lambda e^{x-w}\sqrt{\frac{x}{w^3}}\frac{1}{2}\int_0^\infty e^{-\frac{z}{2}-\frac{(x\vee w)^2+(x\wedge w)^2}{2z}}I_v((x\vee w)(x\wedge w)/z)\frac{dz}{z}dw \\
&= \lambda e^{x-w}\sqrt{\frac{x}{w^3}}\int_0^\infty e^{-\frac{z}{2}-\frac{x^2+w^2}{2z}}I_v(xw/z)\frac{dz}{z}dw,
\end{aligned}$$

where in the third equality we used the identity

$$I_v(x)K_v(w) = \frac{1}{2}\int_0^\infty e^{-\frac{z}{2}-\frac{x^2+w^2}{2z}}I_v(xw/z)\frac{dz}{z}$$

for $w \geq x > 0$ (see (2.7) in [12]). To continue we use another identity for modified Bessel functions (see (2.10) in [12])

$$I_v(r) = \int_0^\infty e^{-\frac{v^2}{2}t}\Theta(r,t)dt,\ r > 0$$

and obtain

$$\begin{aligned}
&\lambda e^{x-w}\sqrt{\frac{x}{w^3}}\int_0^\infty e^{-\frac{z}{2}-\frac{x^2+w^2}{2z}}I_v(xw/z)\frac{dz}{z} \\
&\quad= \lambda e^{x-w}\sqrt{\frac{x}{w^3}}\int_0^\infty \frac{1}{z}e^{-\frac{z}{2}-\frac{x^2+w^2}{2z}}\int_0^\infty e^{-\frac{v^2}{2}t}\Theta(xw/z,t)dt \\
&\quad= \lambda e^{x-w}\sqrt{\frac{x}{w^3}}\int_0^\infty\int_0^\infty e^{-\lambda t-\frac{t}{8}}\Theta(xw/z,t)\frac{1}{z}e^{-\frac{z}{2}-\frac{x^2+w^2}{2z}}dzdt,
\end{aligned}$$

where in the last equality we used Fubini's theorem and that $v^2 = 2\lambda + 1/4$. To finish the proof we compare the last expression with (23). □

Now we consider a family of processes defined by (18) for all $x > 0$, and to underline the dependence on x we denote this process by $\theta(x)$, i.e.

$$\theta_t(x) = \frac{xe^{B_t - \frac{t}{2}}}{1 + x\int_0^t e^{B_u - \frac{u}{2}} du}.$$

From Lemma 3.1 we can deduce

Proposition 3.3. *Let $T_\lambda, \hat{T}_2$ be two independent exponential random variables, which are independent of the Brownian motion B. Then on $(0, \infty)$*

$$z^2 \mathbb{P}\Big(\theta_{T_\lambda}(\hat{T}_2) \in dz\Big) = \mathbb{P}(\hat{T}_2 \in dz)\mathbb{E}(\theta_{T_\lambda}(z))^2. \tag{24}$$

Proof. From Lemma 3.1 for $x > 0$, we have on $(0, \infty)$

$$e^{-2x} z^2 \mathbb{P}(\theta_{T_\lambda}(x) \in dz) = 2\lambda e^{-x-z}\sqrt{xz} F_v(x, z)dz.$$

Thus

$$e^{-2x}\mathbb{E}(\theta_{T_\lambda}(x))^2 = \int_0^\infty 2\lambda e^{-x-z}\sqrt{xz} F_v(x, z)dz.$$

From symmetry of F_v, after integrating on x, we obtain

$$\begin{aligned} z^2 \int_0^\infty e^{-2x}\mathbb{P}(\theta_{T_\lambda}(x) \in dz)dx &= \Big(\int_0^\infty 2\lambda e^{-x-z}\sqrt{xz} F_v(x, z)dx\Big)dz \\ &= e^{-2z}\mathbb{E}(\theta_{T_\lambda}(z))^2 dz. \end{aligned}$$

The assertion follows. □

References

[1] Borodin A., Salminen P. *Handbook of Brownian Motion - Facts and Formulae.* Birkhauser (2nd ed.), 2002.

[2] Comtet A., Monthus C., Yor M. *Exponential Functionals of Brownian motion and disordered systems.* J. Appl. Prob. 35, (1998), 255-271.

[3] Donati-Martin C., Yor M. *Some Brownian functionals and their laws.* Ann. Probab. 25, (1997) 1011-1058.

[4] Donati-Martin C., Matsumoto H., Yor M. *Some absolute continuity relationship for certain anticipative transformations of geometric Brownian motion.* Publ. RIMS, Kyoto Univ. 37 (2001), 295-326.

[5] Jakubowski J., Wiśniewolski M. *On some Brownian functionals and their applications to moments in the lognormal stochastic volatility model*, Stud. Math. 219 (2013), 201-224.

[6] Lungu E.; Øksendal B. *Optimal Harvesting from a Population in a Stochastic Crowded Environment* Mathematical Biosciences 145 (1997), 47-75.
[7] Mackevičius V. *On positive approximations of positive diffusions* Liet. matem. rink, 47, spec. nr., 2007, 58–62
[8] Mackevičius V. *Verhulst versus CIR*, 11th International Vilnius Conference on Probability Theory and Mathematical Statistics (2014).
[9] Mansuy R., Yor M. *Aspects of Brownian Motion.* Universitext, Springer-Verlag, 2008.
[10] H. Matsumoto, M. Yor *An analogue of Pitman's 2M - X theorem for exponential Wiener functionals, part II: The role of the generalized Inverse Gaussian laws*, Nagoya Math. J. 162 (2001), 65-86.
[11] Matsumoto H., Yor M. *A Relationship between Brownian motions with opposite drifts via certain enlargements of the Brownian filtration* Osaka J. Math. 38, (2001) 383-398.
[12] Matsumoto H., Yor M. *Exponential functionals of Brownian motion, I, Probability laws at fixed time.* Probab. Surveys 2, (2005) 312-347.
[13] Matsumoto H., Yor M. *Exponential functionals of Brownian motion, II, Some related diffusion processes.* Probab. Surveys 2, (2005) 348-384.
[14] Øksendal B. *Stochastic Differential Equations*, Springer Verlag, 2000.